arXiv:1101.2503v2 [math.GR] 24 Feb 2011

On the Order of the Schur Multiplier of a Pair of Finite p-Groups

Azam Hokmabadi[a], Fahimeh Mohammadzadeh[a], Behrooz Mashayekhy[b,*]

[a]*Department of Mathematics, Faculty of Sciences, Payame Noor University, Iran.*
[b]*Department of Pure Mathematics, Center of Excellence in Analysis on Algebraic Structures, Ferdowsi University of Mashhad, P. O. Box 1159-91775, Mashhad, Iran.*

Abstract

In 1997, G. Ellis defined the Schur multiplier of a pair (G, N) of groups and mentioned that this notion is a useful tool for studying pairs of groups. In this paper we characterize the structure of a pair of finite p-groups (G, N) in terms of the order of the Schur multiplier of (G, N) under some conditions.

Keywords: Pair of groups, Schur multiplier of a pair, Finite p-groups.
2010 MSC: 20E34, 20D15.

---

1. Introduction and Motivation

A definition for the Schur multiplier of a group G is as the abelian group $M(G) = (R \cap F')/[R, F]$ in which F/R is a free presentation of G. In 1956, J.A. Green [5] showed that the order of the Schur multiplier of a finite p-group of order p^n is bounded by $p^{\frac{n(n-1)}{2}}$, and hence equals to $p^{\frac{n(n-1)}{2}-t}$, for some non-negative integer t. Ya.G. Berkovich [1], X. Zhou [11], G. Ellis [4] and P. Niroomand [8, 9] determined the structure of G for $t = 0, 1, 2, 3, 4, 5$ by different methods.

A pair of groups (G, N) is a group G with a normal subgroup N. In 1998, Ellis [3] defined the Schur multiplier of a pair (G, N) of groups to be

---

*Corresponding author

Email addresses: `ahokmabadi@pnu.ac.ir` (Azam Hokmabadi), `F.mohamadzade@gmail.com` (Fahimeh Mohammadzadeh), `bmashf@um.ac.ir` (Behrooz Mashayekhy)

Preprint submitted to X *February 25, 2011*

the abelian group $M(G,N)$ appearing in the natural exact sequence

$$\begin{aligned} H_3(G) &\rightarrow H_3(G/N) \rightarrow M(G,N) \rightarrow M(G) \rightarrow M(G/N) \\ &\rightarrow N/[N,G] \rightarrow (G)^{ab} \rightarrow (G/N)^{ab} \rightarrow 0 \end{aligned}$$

in which $H_3(G)$ is the third homology of G with integer coefficients. He mentioned that this notion is a useful tool for studying pairs of groups. He also gave an upper bound for the order of the Schur multiplier of pairs of finite groups. It is interesting to know for which classes of pairs of groups the structure of the pair (G,N) can be completely described in terms of the order of $M(G,N)$. In 2004, Salemkar, Moghaddam and Saeedi [10] tried to answer to this question for a pair of finite p-groups and proved the following theorem.

Theorem 1. *([10]) Let (G,N) be a pair of groups and K be the complement of N in G, with $|N| = p^n$ and $|K| = p^m$. Then the following statements hold:*
(i) $|M(G,N)| \leq p^{\frac{1}{2}n(2m+n-1)}$;
(ii) If G is abelian, N is elementary abelian and $|M(G,N)| = p^{\frac{1}{2}n(2m+n-1)}$, then G is elementary abelian;
(iii) If the pair (G,N) is non-capable and $|M(G,N)| = p^{\frac{1}{2}n(2m+n-1)-1}$, then $G \cong \mathbf{Z}_{p^2}$.

In this paper we extend the above theorem and characterize the structure of the pair (G,N) of finite p-groups in terms of the order of $M(G,N)$ in more cases. Let G, N and K be as in Theorem 1 and $|M(G,N)| = p^{\frac{1}{2}n(2m+n-1)-t}$. Then we prove $t = 0$ if and only if N is trivial or (G,N) is a pair of elementary abelian p-groups. Also, we determine the pair (G,N), for $t = 1$. Moreover, we give the structure of (G,N) for $t = 2$ and $t = 3$, when K is a normal subgroup of G. The main results of this paper are somehow generalizations of the results of [1], [11] and [4] to the pair of finite p-groups

2. Preliminaries

Let (G,N) be a pair of groups. We recall that a relative central extension of the pair (G,N) consists of a group homomorphism $\sigma : M \rightarrow G$, together with an action of G on M, such that
(i) $\sigma(M) = N$,
(ii) $\sigma(m^g) = g^{-1}\sigma(m)g$, for all $g \in G$, $m \in M$
(iii) $m^{\sigma(m_1)} = m_1^{-1}mm_1$, for all $m, m_1 \in M$

(iv) G acts trivially on $\ker \sigma$.

The G-commutator subgroup of M is defined to be the subgroup $[M, G]$ generated by the G-commutators $[m, g] = m^{-1}m^g$ for all $g \in G, m \in M$ and the G-center of M is the central subgroup

$$Z(M, G) = \{m \in M | m^g = m \ \ for\ all\ g \in G\}.$$

Also, the subgroup $Z_2(M, G)$ is defined by

$$\frac{Z_2(M, G)}{Z(M, G)} = Z(\frac{M}{Z(M, G)}, G).$$

A pair (G, N) is said to be capable if it admits a relative central extension $\sigma : M \to G$ with $\ker \sigma = Z(M, G)$. Note that a group G is capable precisely when the pair (G, G) is capable.

We call a pair (G, N) an extra special pair of p-groups when $Z(N, G)$ and $[N, G]$ are the same subgroups of order p. Also, we need to recall the definition of a covering pair.

Definition 2. *([3]) A relative central extension $\sigma : N^* \to G$ of the pair (G, N) will be called a covering pair if there exists a subgroup A of N^* such that*
(i) $A \leq Z(N^*, G) \cap [N^*, G]$;
(ii) $A \cong M(G, N)$;
(iii) $N \cong N^*/A$.

The following theorem plays an important role to prove the main results.

Theorem 3. *([3]) Let (G, N) be a pair of groups and K be the complement of N in G. Then*

$$M(G) \cong M(G, N) \times M(K).$$

We recall from [6] that if $G = N \times K$, then

$$M(G) \cong M(N) \times M(K) \times (N^{ab} \otimes K^{ab}).$$

A useful consequence of this fact is as follows.

Corollary 4. *If $G = N \times K$, then*

$$|M(G, N)| = |M(N)||N^{ab} \otimes K^{ab}|.$$

We use the following theorems in the next section. Here D denotes the dihedral group of order 8, Q denotes the quaternion group of order 8 and E_1 and E_2 denote the extra special groups of order p^3 with odd exponent p and p^2, respectively.

Theorem 5. *([4]) Let G be a group of prime-power order p^n. Suppose that $|M(G)| = p^{\frac{1}{2}n(n-1)-t}$. Then*
(i) $t = 0$ if and only if G is elementary abelian ([1]);
(ii) $t = 1$ if and only if $G \cong \mathbf{Z}_{p^2}$ or $G \cong E_1$ ([1]);
(iii) $t = 2$ if and only if $G \cong \mathbf{Z}_p \times \mathbf{Z}_{p^2}$, $G \cong D$ or $G \cong \mathbf{Z}_p \times E_1$ ([11]);
(iv) $t = 3$ if and only if $G \cong \mathbf{Z}_{p^3}$, $G \cong \mathbf{Z}_p \times \mathbf{Z}_p \times \mathbf{Z}_{p^2}$, $G \cong Q$, $G \cong E_2$, $G \cong D \times \mathbf{Z}_2$ or $G \cong E_1 \times \mathbf{Z}_p \times \mathbf{Z}_p$.

Theorem 6. *([6]) Let $G \cong \mathbf{Z}_{n_1} \times \mathbf{Z}_{n_2} \times ... \times \mathbf{Z}_{n_k}$, where $n_{i+1}|n_i$ for all $i \in 1, 2, ..., k-1$ and $k \geq 2$, and let $\mathbf{Z}_n^{(m)}$ denote the direct product of m copies of $\mathbf{Z}_n$. Then*

$$M(G) \cong \mathbf{Z}_{n_2} \times \mathbf{Z}_{n_3}^{(2)} \times ... \times \mathbf{Z}_{n_k}^{(k-1)}.$$

Theorem 7. *([3]) Let (G, N) be a pair of groups such that $N/Z(N, G)$ is finite of prime power order p^n and G/N is finite of prime power order p^m. Then*

$$|M(G, N)||[N, G]| \leq p^{n(2m+n-1)/2}.$$

Theorem 8. *([10]) Let (G, N) be a pair of finite p-groups with G/N and $N/Z(N, G)$ of orders p^m and p^n, respectively. If $|[N, G]| = p^{\frac{1}{2}n(2m+n-1)}$, then either $N/Z(N, G)$ is elementary abelian or the pair $(G/Z(N, G), N/Z(N, G))$ is an extra special pair of finite p-groups.*

Theorem 9. *([10]) Let (G, M) be a pair of groups with G/M and $M/Z(M, G)$ of orders p^m and p^n, respectively. Suppose $z \in Z_2(M, G) - Z(M, G)$ and consider two non-negative integers $\mu(z)$ and $\nu(z)$ where*

$$p^{\mu(z)} = |[G, z]| \quad , p^{\nu(z)} = |\frac{G/[G, z]}{Z(G/[G, z]), M/[G, z])}|.$$

Then
(a) $|[M, G]| \leq p^{\frac{1}{2}(\nu(z)(\nu(z)-1)-m(m-1))+\mu(z)} \leq p^{\frac{1}{2}n(2m+n-1)}$.
(b) Suppose for some none-negative integer s, $|[M, G]| = p^{n(2m+n-1)/2-s}$, then

the following statements hold:
(i) $|[M/Z(M,G), G/Z(M,G)]| \leq p^{s+1}$. *If* $|[M/Z(M,G), G/Z(M,G)]| = p^{s+1-k}$ *for some* $0 \leq k \leq s+1$, *then* $exp(Z_2(M,G)/Z(M,G)) \leq p^{k+1}$ *and* $m+n-1-s \leq \mu(z) \leq m+n-1-s+k$.
(ii) If $exp(Z_2(M,G)/Z(M,G)) \geq p^k$, *then* $m+n \leq s/(k-1)+k/2$.

3. Main Results

In this section we always assume that (G,N) is a pair of finite p-groups such that K is the complement of N in G, with $|N| = p^n$ and $|K| = p^m$, and hence $|M(G,N)| = p^{\frac{1}{2}n(2m+n-1)-t}$, for some $t \geq 0$.

Salemkar, Moghaddam and Saeedi [10] proved that if $t = 0$ and G is abelian and N is elementary abelian, then G is elementary abelian. The first main result of this paper gives a suitable extended version of the above result similar to Berkovich's one [1].

Theorem 10. *With the previous assumptions and notation,* $t = 0$ *if and only if* N *is trivial or* (G,N) *is a pair of elementary abelian* p*-groups.*

Proof. Using Theorem 3, we have $|M(G,N)| = |M(G)|/|M(K)|$. Hence necessity is immediate by Theorems 5. For sufficiency let the relative central extension $\sigma : N^* \to G$ be a covering pair of (G,N). Then there exists a subgroup A of N^* such that $A \leq Z(N^*,G) \cap [N^*,G]$, $A \cong M(G,N)$ and $N \cong N^*/A$. It is easy to see that for any $k \in K$, the map $\varphi_k : N^* \to N^*$, defined by $\varphi_k(n^*) = n^{*k}$, is an automorphism of N^*. Therefore, using the homomorphism $\psi : K \to Aut(N^*)$, given by $\psi(k) = \varphi_k$, we can define a semidirect product of N^* by K, denoted by G^*. It is straightforward to check that the subgroups $[N^*,G]$ and $Z(N^*,G)$ are contained in $[N^*,G^*]$ and $Z(N^*,G^*)$ respectively. Then the map $\delta : G^* \to G$, given by $\delta(n^*k) = \sigma(n^*)k$, for $n^* \in N^*$, $k \in K$, is an epimorphism with $\ker(\delta) = \ker(\sigma)$. Therefore

$$|\frac{N^*}{Z(N^*,G^*)}| \leq |\frac{N^*}{Z(N^*,G)}| \leq |\frac{N^*}{A}| = |N| = p^n$$

and

$$|\frac{G^*}{N^*}| = |\frac{G^*/A}{N^*/A}| = |\frac{G}{N}| = |K| = p^m.$$

So $|[N^*,G^*]| \leq p^{\frac{1}{2}n(2m+n-1)}$ by Theorem 7. This implies that

$$p^{\frac{1}{2}n(2m+n-1)} = |M(G,N)| = |A| \leq |[N^*,G^*]| \leq p^{\frac{1}{2}n(2m+n-1)}.$$

Thus $A = [N^*, G^*]$. It follows that $N \leq Z(G)$ and we have $G = N \times K$. Then Corollary 4 implies that $p^{\frac{1}{2}n(2m+n-1)} \leq p^{\ \frac{1}{2}n(n-1)}|N^{ab} \otimes K^{ab}|$. Hence $p^{mn} \leq |N^{ab} \otimes K^{ab}| \leq p^{md(N)}$, where $d(N)$ is the minimum number of generators of N. Therefore $n = d(N)$ and hence N is an elementary abelian p-group and $|N^{ab} \otimes K^{ab}| = p^{mn}$. If $n = 0$, then N is trivial subgroup. If $n > 0$, then we have $p^{nm} = |\mathbf{Z}_p^{(n)} \otimes K^{ab}| = p^{d(K^{ab})n}$ and it follows that $d(K^{ab}) = m$. Therefore $d(K) = m$ and so K is an elementary abelian p-group too. This completes the proof. □

Lemma 11. *Let* (G, N) *be a pair of* p*-groups such that* $[N, G] \neq 1$. *Then* $Z(N, G) \cap [N, G] \neq 1$.

Proof. Using the fact that $Z(N, G) = Z(G) \cap N$, the result follows. □

Salemkar, Moghaddam and Saeedi [10] proved that if $t = 1$ and (G, N) is non-capable, then $G \cong \mathbf{Z}_{p^2}$. The second main result of this paper gives a vast generalization of the above result similar to Berkovich's one [1].

Theorem 12. *With the previous assumptions if* $t = 1$, *then one of the following cases holds:*
(i) $G \cong N \times K$ *where* $N \cong \mathbf{Z}_{p^2}$ *and* $K = 1$*;*
(ii) $G \cong N \times K$, *where* $N \cong \mathbf{Z}_p$ *and* K *is any group with* $d(K) = m - 1$*;*
(iii) (G, N) *is an extra special pair of groups which is capable.*

Proof. Choose N^*, G^* and A as in the proof of Theorem 10. We divide the proof in two cases:
Case 1. Suppose $A \neq Z(N^*, G^*)$. Then $|N^*/Z(N^*, G^*)| < |N^*/A| = p^n$ and $|N^*/Z(N^*, G^*)| \leq p^{n-1}$. Thus by Theorem 7

$$p^{\frac{1}{2}n(2m+n-1)-1} = |A| \leq |[N^*, G^*]| \leq p^{\frac{1}{2}(n-1)(2m+n-2)}.$$

This implies that $n + m \leq 2$. Since $t = 1$, by Theorem 10 we have $n \neq 0$ and G is not an elementary abelian p-group. Therefore we have $G = N \cong \mathbf{Z}_{p^2}$.
Case 2. Suppose $A = Z(N^*, G^*)$. Then $|N^*/Z(G^*, N^*)| = p^n$. By Theorem 7, we have

$$p^{\frac{1}{2}n(2m+n-1)-1} = |A| \leq |[N^*, G^*]| = p^{\frac{1}{2}n(2m+n-1)-s}$$

for some $s \geq 0$. It follows that $s \leq 1$.

First, assume that $s = 0$. Hence Theorem 9 implies that $|[N, G]| = p$ and $\exp(Z(N, G)) = p$. If $Z(N, G)$ is cyclic, then $[N, G] = Z(N, G)$ has order p by Lemma 11 and hence (G, N) is an extra special pair of groups which is capable. If $Z(N, G)$ is not cyclic, then $Z(N, G) = Z_2(N^*, G^*)/Z(N^*, G^*)$ has two distinct subgroups of order p. Therefore there exist elements $y_0, z_0 \in Z_2(N^*, G^*) - Z(N^*, G^*)$ such that

$$| < y_0 Z(N^*, G^*) > | = | < z_0 Z(N^*, G^*) > | = p$$

and $< y_0 Z(N^*, G^*) > \cap < z_0 Z(N^*, G^*) >= 1$. Using the proof of Theorem 9, we have $|[G^*, y_0]| = |[G^*, z_0]| = p^{m+n-1}$. On the other hand, $[G^*, y_0] \cong G/C_{G^*}(y_0)$ and $[G^*, z_0] \cong G/C_{G^*}(z_0)$. So $|C_{G^*}(y_0)| = |C_{G^*}(z_0)| = p$. Thus we have

$$Z(N^*, G^*) \leq C_{G^*}(y_0) \cap C_{G^*}(z_0) = [N^*, G^*].$$

This implies that $|[N, G]| = |[N^*/Z(G^*, N^*), G^*/Z(G^*, N^*)]| = 1$ which is a contradiction.

Now assume that $s = 1$, then $A = [N^*, G^*]$. This implies that $N \leq Z(G)$. Thus $G \cong N \times K$. Then by Corollary 4 we have $|M(G, N)| = |M(N)||N^{ab} \otimes K^{ab}|$. Thus $p^{\frac{1}{2}n(2m+n-1)-1} \leq p^{\frac{1}{2}n(n-1)} \times |N^{ab} \otimes K^{ab}|$. Hence $p^{mn-1} \leq |N^{ab} \otimes K^{ab}| \leq p^{md(N)}$. Therefore $m(n - d(N)) \leq 1$. Then $m = 0$ or $n - d(N) = 0$ or $m = n - d(N) = 1$. If $m = 0$, then $G = N \cong \mathbf{Z}_{p^2}$ by Theorem 5. If $d(N) = n$, then $N \cong \mathbf{Z}_p^{(n)}$. Hence $p^{d(K)n} = |N^{ab} \otimes K^{ab}| = p^{mn-1}$ and $n(m - d(K)) = 1$. Therefore $n = 1$ and $d(K) = m - 1$. In other words, $N \cong \mathbf{Z}_p$ and K is any group with $d(K) = m-1$. Finally, if $m = n-d(N) = 1$, then $K \cong \mathbf{Z}_p$ and $N \cong \mathbf{Z}_{p^2} \times \mathbf{Z}_p^{(n-2)}$ which by Corollary 4 and Theorem 6 we must have $n = 1$ which is a contradiction. □

With an additional condition we will be able to state the reverse of the above theorem as follows.

Theorem 13. *Let K be a normal subgroup of G. Then $t = 1$ if and only if $G \cong N \times K$ with one of the following cases:*
i) $N \cong \mathbf{Z}_{p^2}$ and $K = 1$;
ii) $N \cong \mathbf{Z}_p$ and K is any group with $d(K) = m - 1$;
iii) $N \cong E_1$ and $K = 1$.

Proof. By the assumption K is normal and so $G \cong N \times K$. Hence by Corollary 4 we have

$$|M(G, N)| = |M(N)||N^{ab} \otimes K^{ab}|. \tag{1}$$

Necessity is clear by the above equality. For sufficiency, first we suppose that N is an elementary abelian p-group. Then $N \cong \mathbf{Z}_{p^n}$ and $|N^{ab} \otimes K^{ab}| = |M(G,N)|/|M(N)| = p^{mn-1}$. On the other hand we have $|N^{ab} \otimes K^{ab}| = |\mathbf{Z}_p^{(n)} \otimes K^{ab}| = p^{nd(K)}$. Therefore $mn - 1 = nd(K)$ and so $n(m - d(K)) = 1$. This implies that $N \cong \mathbf{Z}_p$ and $d(K) = m - 1$.

Now suppose that N is not elementary abelian. Then using (1) we have

$$|M(G,N)| < p^{\frac{1}{2}n(n-1)}|K|^{d(N)} \leq p^{\frac{1}{2}n(n-1)+md(N)}.$$

Hence $t = 1$ implies that $m(n - d(N)) = 0$. But $n \neq d(N)$. Therefore $m = 0$ and $|M(G,N)| = |M(N)| = p^{\frac{1}{2}n(n-1)-1}$. Hence by Theorem 5 we have $K = 1$ and $N \cong \mathbf{Z}_{p^2}$ or $N \cong E_1$. This completes the proof. $\square$

Our third main result is somehow a generalization of the Zhou's one [11] to the pair of finite p-groups.

Theorem 14. *Let K be a normal subgroup of G. Then $t = 2$ if and only if $G \cong N \times K$ with one of the following cases:*
(i) $N \cong \mathbf{Z}_p \times \mathbf{Z}_{p^2}$ and $K = 1$;
(ii) $N \cong D$ and $K = 1$;
(iii) $N \cong \mathbf{Z}_p \times E_1$ and $K = 1$;
(iv) $N \cong \mathbf{Z}_{p^2}$ and $K \cong \mathbf{Z}_p$;
(v) $N \cong E_1$ and $K \cong \mathbf{Z}_p$;
(vi) $N \cong \mathbf{Z}_p \times \mathbf{Z}_p$ and K is any group with $d(K) = m - 1$;
(vii) $N \cong \mathbf{Z}_p$ and K is any group with $d(K) = m - 2$.

Proof. Since K is a normal subgroup, we have $G \cong N \times K$. Hence by Corollary 4, (1) holds and necessity follows. For sufficiency, we proceed as in Theorem 13. First suppose that N is an elementary abelian p-group. Then $|N^{ab} \otimes K^{ab}| = |M(G,N)|/|M(N)| = p^{mn-2}$. On the other hand, we have $|N^{ab} \otimes K^{ab}| = p^{nd(K)}$. Hence $mn - 2 = nd(K)$ and so $n(m - d(K)) = 2$. It follows that $n = 1$ or $n = 2$. If $n = 1$, then $N \cong \mathbf{Z}_p$ and K is any group with $d(K) = m - 2$. If $n = 2$, then $N \cong \mathbf{Z}_p \times \mathbf{Z}_p$ and K is any group with $d(K) = m - 1$.

Now suppose that N is not an elementary abelian p-group. Then (1) implies that $|M(G,N)| < p^{\frac{1}{2}n(n-1)}|K|^{d(N)} \leq p^{\frac{1}{2}n(n-1)+md(N)}$. It follows that $m(n - d(N)) < 2$. Therefore $m = 0$ or $m = 1$. If $m = 0$, then by Theorem 5 we have $K = 1$ and $N \cong \mathbf{Z}_p \times \mathbf{Z}_{p^2}$ or $N \cong D$ or $N \cong \mathbf{Z}_p \times E_1$. If $m = 1$ then $K \cong \mathbf{Z}_p$ and $d(N) = n-1$. So $|N^{ab} \otimes K^{ab}| = p^{n-1}$. Therefore the equality (1) implies that $|M(N)| = p^{\frac{1}{2}n(n-1)-1}$. Now using Theorem 5 we have $N \cong \mathbf{Z}_{p^2}$ or $N \cong E_1$. This completes the proof. $\square$

Finally, our last main result is somehow a generalization of the Ellis' one [4] to the pair of finite p-groups.

Theorem 15. *Let* K *be a normal subgroup of* G*. Then* $t = 3$ *if and only if* $G \cong N \times K$ *with one of the following cases:*
(i) $N \cong \mathbf{Z}_{p^3}$ *and* $K = 1$*;*
(ii) $N \cong \mathbf{Z}_p \times \mathbf{Z}_p \times \mathbf{Z}_{p^2}$ *and* $K = 1$*;*
(iii) $N \cong Q$ *and* $K = 1$*;*
(iv) $N \cong E_2$ *and* $K = 1$*;*
(v) $N \cong D \times \mathbf{Z}_2$ *and* $K = 1$*;*
(vi) $N \cong E_1 \times \mathbf{Z}_p \times \mathbf{Z}_p$ *and* $K = 1$*;*
(vii) $N = K \cong \mathbf{Z}_{p^2}$*;*
(viii) $N \cong \mathbf{Z}_{p^2} \times \mathbf{Z}_p$ *and* $K \cong \mathbf{Z}_p$*;*
(ix) $N \cong D$ *and* $K \cong \mathbf{Z}_p$*;*
(x) $N \cong \mathbf{Z}_p \times E_1$ *and* $K \cong \mathbf{Z}_p$*;*
(xi) $N \cong \mathbf{Z}_{p^2}$ *and* $K \cong \mathbf{Z}_p \times \mathbf{Z}_p$*;*
(xii) $N \cong E_1$ *and* $K \cong \mathbf{Z}_p \times \mathbf{Z}_p$*;*
(xiii) $N \cong \mathbf{Z}_p$ *and* K *is any group with* $d(K) = m - 3$*;*
(xiv) $N \cong \mathbf{Z}_p \times \mathbf{Z}_p \times \mathbf{Z}_p$ *and* K *is any group with* $d(K) = m - 1$*.*

Proof. Necessity is straightforward. The proof of sufficiency is similar to the proof of previous theorems. Suppose that N is an elementary abelian p-group. Since $t = 3$, we have $p^{nd(K)} = |N^{ab} \otimes K^{ab}| = p^{mn-3}$. This implies that $n(m - d(K)) = 3$. So $n = 1$ or $n = 3$. If $n = 1$ then $N \cong \mathbf{Z}_p$ and K is any group with $d(K) = m - 3$. If $n = 3$ then $N \cong \mathbf{Z}_p \times \mathbf{Z}_p \times \mathbf{Z}_p$ and K is any group with $d(K) = m - 1$.

Now suppose that N is not an elementary abelian p-group. Then we have $|M(G, N)| < p^{\frac{1}{2}n(n-1)+md(N)}$ and so $m(n - d(N)) \leq 2$. This implies that $m = 0$, $m = 1$ or $m = 2$.

If $m = 0$, then by Theorem 5 we have $K = 1$ and $N \cong \mathbf{Z}_{p^3}$ or $N \cong \mathbf{Z}_p \times \mathbf{Z}_p \times \mathbf{Z}_{p^2}$ or $N \cong Q$ or $N \cong E_2$ or $N \cong D \times \mathbf{Z}_2$ or $N \cong E_1 \times \mathbf{Z}_p \times \mathbf{Z}_p$.

If $m = 1$, then $K \cong \mathbf{Z}_p$ and $d(N) = n - 1$ or $d(N) = n - 2$. If $d(N) = n - 1$, then $|N^{ab} \otimes K^{ab}| = p^{n-1}$ and so $|M(N)| = |M(G, N)|/|N^{ab} \otimes K^{ab}| = p^{\frac{1}{2}n(n-1)-2}$. It follows that $N \cong \mathbf{Z}_{p^2} \times \mathbf{Z}_p$ or $N \cong D$ or $N \cong \mathbf{Z}_p \times E_1$, by Theorem 5. If $d(N) = n - 2$, then $|M(N)| = p^{\frac{1}{2}n(n-1)-1}$ and hence $N \cong \mathbf{Z}_{p^2}$ or $N \cong E_1$ which is a contradiction.

If $m = 2$, then $d(N) = n-1$ and $K \cong \mathbf{Z}_{p^2}$ or $K \cong \mathbf{Z}_p \times \mathbf{Z}_p$. If $K \cong \mathbf{Z}_p \times \mathbf{Z}_p$, then $|N^{ab} \otimes K^{ab}| = p^{2(n-1)}$ and so $|M(N)| = p^{\frac{1}{2}n(n-1)-1}$. Then $N \cong \mathbf{Z}_{p^2}$ or $N \cong E_1$. Now suppose $K \cong \mathbf{Z}_{p^2}$. Then $N^{ab} \cong \mathbf{Z}_p^{(n-1)}$ or $N^{ab} \cong \mathbf{Z}_{p^2} \times \mathbf{Z}_p^{(n-2)}$. If $N^{ab} \cong \mathbf{Z}_p^{(n-1)}$, then $|N^{ab} \otimes K^{ab}| = p^{n-1}$ and so $|M(N)| = p^{\frac{1}{2}(n^2+n-4)}$ which is a contradiction. If $N^{ab} \cong \mathbf{Z}_{p^2} \times \mathbf{Z}_p^{(n-2)}$, then $|N^{ab} \otimes K^{ab}| = p^n$ and hence $|M(N)| = p^{\frac{1}{2}(n^2+n-6)}$ which implies that $n \leq 2$. Therefore $N \cong \mathbf{Z}_{p^2}$. This completes the proof. $\square$

References

[1] Ya. G. Berkovich, On the order of the commutator subgroup and the Schur multiplier of a finite p-group, J. Algebra 144 (1991) 269-272.

[2] G. Ellis, Capability, homology, and central series of a pair of groups, J. Algebra 179 (1995) 31-46.

[3] G.Ellis, The Schur multiplier of a pair of groups, Appl. Categ. Structures 6 (1998) 355-371.

[4] G.Ellis, On the Schur multiplier of p-groups, Comm. Algebra 27 (9) (1999) 4173-4178.

[5] J.A. Green, On the number of outomorphisms of a finite p-group, Proc. Roy. Soc. London Ser. A 237 (1958) 574-581.

[6] G. Karpilovsky, The Schur Multiplier, London Math. Soc. Monographs, New Series 2, 1987.

[7] J.-L. Loday, Cohomologie et group de Steinberg relatif, J. Algebra 54 (1978) 178-202.

[8] P. Niroomand, Characterizing finite p-groups by their Schur multipliers, available at arXiv: math. Gr/ 1001.4256.

[9] P. Niroomand, Characterizing finite p-groups by their Schur multipliers, $t(G) = 5$, available at arXiv: math. Gr/ 1001.4257.

[10] A. Salemkar, M. R. Moghaddam and F. Saeedi, The commutator subgroup and Schur multiplier of finite p-groups , J. Aust. Math. Soc. 81 (2006) 1-9.

[11] X. Zhou, On the order of Schur multipliers of finite p-groups, Comm. Algebra 22 (1) (1994) 1-8.